

**European Project
AIR3CT93-0818**

VEGEXTRACT

Fátima Rodrigues
Gabriela Bernardo-Gil
José Abecassis Empis
Michel Rouzet
Regina Santos
Tiejun Lu
Michael King
João Bastos

**A contribution to the study of
Rosmarinus officinalis L.
from Portugal**



Ed:
Gabriela Bernardo-Gil
José Abecassis Empis



**AIR
AGRO-INDUSTRIAL RESEARCH**

European Commission - DG XII - Science, Research and Development

TABLE OF CONTENTS

1. INTRODUCTION

- 1.1. *Rosmarinus officinalis* and Industry
- 1.2. Traditional Uses
- 1.3. Official Standards

2. BOTANICAL DESCRIPTION

- 2.1. Morphologic characteristics
- 2.2. Classification

3. VEGETABLE MATERIAL STUDIED

- 3.1. Source and pre-treatment of the plants used in the present study
- 3.2. Choice of the best period to harvest the plant
 - 3.2.1. Methodology
 - 3.2.2. Yield of the aromatic extract
 - 3.2.3. Alteration of the chemical composition of the aromatic extract

4. EXTRACTION OF AROMATIC PRODUCTS FROM ROSEMARY

- 4.1. Extraction of flavour components
 - 4.1.1. Production of essential oil
 - 4.1.2. Extraction with organic solvents
 - 4.1.3. Extraction with dense gases
 - 4.1.4. Recent developments in extraction
- 4.2. Enzymatic pre-treatment

5. YIELD OF AROMATIC EXTRACT

- 5.1. Distillation methods
- 5.2. Extraction with supercritical fluids

6. CHEMICAL ANALYSIS OF THE DIFFERENT EXTRACTS FROM *ROSMARINUS OFFICINALIS*

- 6.1. Generalities
- 6.2. Chromatographic study of the essential oils and CO₂ extracts
 - 6.2.1. Methodology used
 - 6.2.2. Composition of the essential oil
- 6.3. Study of the extract obtained by extraction with supercritical and marginally sub-critical fluids
 - 6.3.1. Extraction with liquid CO₂
 - 6.3.2. Extraction with supercritical CO₂

7. ANTIOXIDANTS FROM ROSEMARY

- 7.1. Interest of antioxidants from herbs
- 7.2. Antioxidant extraction
 - 7.2.1. Traditional extraction
 - 7.2.2. Extraction with supercritical fluids
- 7.3. Evaluation of antioxidant capacity
 - 7.3.1. Classical methods
 - 7.3.2. Rancimat®
- 7.4. Yield of antioxidant extracts
- 7.5. Determination of AO capacity of the extracts
- 7.6. Conclusions

8. SUPERCRITICAL EXTRACTION DATA MODELLING

- 8.1. Introduction
- 8.2. Extraction Models
 - 8.2.1. Single Sphere Model I

8.2.2. Single Sphere Model II

8.2.3 Characteristic time model

8.3. Data Modelling Results

8.3.1. Modelling of experimental data obtained with Belmonte's flowered rosemary at 20 MPa, 313 K, and CO₂ flow 5.9 gmin⁻¹, in Bm.

8.3.2. Modelling of experimental data obtained with flowered rosemary from Belmonte at 10 MPa, 313 K, and CO₂ flow 3.33 gmin⁻¹, in Lx.

8.3.3. Modelling of experimental data obtained with flowered rosemary from Belmonte at 10 MPa, 313 K, and CO₂ flow 3.33 gmin⁻¹, in Lx, with total extraction conditions.

8.3.4. Modelling of experimental data obtained with flowered rosemary from Belmonte at 10 MPa, 298 K, and CO₂ flow 3.05 gmin⁻¹, in Bm.

8.3.5. Modelling of experimental data obtained with flowered rosemary from Belmonte at 10 MPa, 298 K, and CO₂ flow 11.30 gmin⁻¹, in Bm.

8.3.6. Modelling of experimental data obtained with flowered rosemary from Belmonte at 20 MPa, 313 K, and CO₂ flow 5.9 gmin⁻¹, in Bm.

8.3.7. Modelling of experimental data obtained with flowered rosemary from Belmonte at 10 MPa, 313 K, and CO₂ flow 5.9 gmin⁻¹, in Bm.

8.3.8. Modelling of experimental data obtained with flowered rosemary from Belmonte at 25 MPa, 313 K, and CO₂ flow 5. gmin⁻¹, in Bm.

8.4. Data Modelling Discussion

9. CONCLUSIONS

10. REFERENCES